

Docket No.: 0259-0411PUS1  
(PATENT)

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

---

In re Patent Application of:  
Maurizio Dalle CARBONARE et al.

Application No.: 10/019,387

Confirmation No.: 6340

Filed: March 26, 2003

Art Unit: 1612

For: USE OF HYALURONIC ACID DERIVATIVES      Examiner: S. Macwall  
FOR THE PREPARATION OF  
PHARMACEUTICAL COMPOSITIONS AND  
BIOMATERIALS FOR THE PREVENTION OF  
THE FORMATION AND URE OF  
CUTANEOUS SCARS

---

**ABATANGELO DECLARATION - DECLARATION UNDER 37 CFR 1.132**

Commissioner for Patents  
P.O. Box 1450  
Alexandria, VA 22313-1450

Sir:

I, Giovanni Abatangelo, do hereby declare the following:

1. Attached is a copy of my *curriculum vitae*.
2. I am presently a Full Professor of Histology and Embryology at the University of Padova, Italy.
3. I have over 25 years of experience in research on tissue engineering by use of hyaluronic acid based biomaterials.
4. I am familiar with the above referenced patent application, as well as the development, usages and properties of hyaluronic acid derivatives and their uses, in particular to reducing normotrophic scarring.

5. I have read and understand the subject matter of the Office Action of September 04, 2008.
6. I am familiar with the Declaration of Anna Maria Zanellato executed on February 3, 2009, which discussed experimental reports and offered comments in support of the patentability of the instant invention, and which Declaration was submitted to the USPTO with an Amendment on February 3, 2009.
7. I offer the following comments as further explanation of the experimental results submitted with the Zanellato Declaration, and to respond to some questions I understand were raised by the Examiners during an interview on February 25, 2009.
8. Explanation of the test result graph in Attachment 2 – I understand the Examiners questioned the meaning of the results in the graph of Attachment 2 of the prior Zanellato Declaration, namely the graph of the results of the test for “Areas of scarring”. I understand that the Examiners questioned why the areas of scarring for samples A and B are low at day 1, are then about equal to C and D on day 3, and then are again reduced at days 14 and 42.

8.1. The process of wound healing involves three major overlapping phases: lag phase (inflammation and cytokine release), proliferation phase (epithelial cells and fibroblasts migration and proliferation, connective tissue deposition and angiogenesis), and remodeling phase (with cell number decreases and collagen fiber organization). (See Exhibit 1)

8.2. During the initial healing phase encompassing the first three to four days after wounding, the process of wound healing is initiated by thrombogenic stimuli, and immune cells infiltrate the injury site where platelets release a variety of cytokines and growth factors. Therefore, during this phase, inflammatory components prevail.

8.3. The proliferative phase (10 and 14 days after wounding) results in regeneration of epidermis, neoangiogenesis and proliferation of fibroblasts with increased collagen synthesis and closure of the skin defect.

8.4. The final remodeling phase takes place over 6 months.

8.5. In my opinion, it is evident that the areas of scarring for samples A and B are about equal to C and D on day 3 because this is the time of lag phase where inflammatory components prevail over the proliferation phase and, therefore, new tissue is not yet made. On day 1, the lag phase is starting and one can value an initial effect of the samples according to the invention. Nevertheless, on day 3 the lag phase is to maximum expression and the inflammation prevails over the samples' effects. Only in the proliferation phase (on day 14), can one estimate the real effect of the samples A and B according to the present invention.

9. Explanation of test results in Attachment 1 – I also understand that the Examiners requested some further explanation of the graph results in Attachment 1, namely the graph relating to “Effective hyaluronate formulations on wound coverage”. Indication in the graph of the “wound coverage” is based on a score related to the length of epidermal coverage as a measure of “re-epithelialization”. Thus, an increased number for “wound coverage” represents an increased amount of re-epithelialization, which is a positive result. The graph shows that tests D and E according to the present invention exhibited increased wound coverage (namely improved re-epithelialization); whereas, the comparative tests A, B and C showed wound coverage values either about the same as control or actually less than control.

In addition, I understand that the data presented below in Table 1 were provided in a newly submitted Zanellato Declaration 2, which the quantitative values of improvement for the tested samples are provided, wherein samples D and E according to the present invention were

compared to the other samples, and wherein it is assumed that the value of wound coverage obtained with the control "alginate vehicle" is 100.


Table 1:

A	B	C	D	E	F
100	110	101	140	138	110

From these results it can be seen that samples D and E according to the present invention exhibited an improvement of about 40% in the wound coverage versus all control samples.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Dated: March 9<sup>th</sup>, 2009

By   
Giovanni Abatangelo

1

*Prof. Abatangelo*  
*Dept. Histology, Microbiology and Biomedical Technology*  
*University of Padova, Italy*

Full Professor of Histology and Embriology, University of Padova, Italy.

Graduated as MD (University of Padua, 1965), Postdoctoral Specialization in Clinical Pathology (University of Padua, 1977), Postdoctoral Fellow Dept. of Biochemistry, Baylor University of Baylor, Houston, Texas, Usa, (1965-1967), Lecturer, Institute of Histology, University of Padua (1970), Full Professor of Histology - Embriology, University of Padua (1975 - present).

President of the Italian Society of Cutaneous Biology and Member of the Editorial Board of "Wound Repair and Regeneration" (Mosby Inc., St. Louis MO, USA).

Organizer and President of the following Symposia:

- 1° and 2° International Symposium on Cutaneous Development , Aging and Repair, Padua, Oct. 4-7 1987 and Oct. 13-16 1991.

- 5° Annual Meeting of the European Tissue Repair Society.

President of the Consortium "TissueTech", a non-profit joint collaborative effort between the University of Padua and Biotechnological Industries dedicated to research project in biomaterials and tissue engineering for clinical use supported by the Italian Government, art. 10, L. 46/82(1989-2001).

Coordinator of research programs that obtained grants from Italian Ministry of University and Research (60%) and Italian National Council Research for the research project "Biomaterials in Orthopedics: Scaffolds for Bioengineered Cartilage".

Research Leader of Four-Year Grant from the European Community (Number QLK3-CT-1999-00625) entitled "Peripheral Nerve Repair"

Author and co-author of 60 papers on peer reviewed journals. During the last decade the interest is mainly devoted to tissue engineering of skin, cartilage and guided regeneration of small arteries.

Prof. Abatangelo research activities have focused for many years on tissue engineering by means hyaluronic acid-derived biomaterials as scaffold for the creation of in vitro artificial tissue such as: skin, cartilage, bone. "Smart scaffold" biomaterials are comprised of hyaluronan derivatives, namely the benzyl ester of this extracellular polysaccharide. All these scaffolds are highly biocompatible and when placed inside the human body, do not elicit any adverse reactions and are resorbed by the host tissues. Abatangelo's laboratories have cultured different kinds of cellular types:

- human dermal fibroblasts: able to attach to the biomaterial fibers, proliferate, colonize the entire device, and secrete the main extracellular molecules (collagen type I, III and IV, laminin, fibronectin) giving rise to a dermal-like tissue;
- keratinocytes able to give rise to a complete skin equivalent when they are in vitro cultured onto this dermal substitute [Zacchi 1998; Galassi 2000; Tonello 2005],
- endothelial cells seeded into this in vitro reconstructed connective tissue are able to proliferate and organize themselves into microcapillary structures [Tonello 2003];
- chondrocytes were cultured inside non-woven Hyaff™ (Hyaluronic benzilic ester, FIDIA Abano Terme) scaffolds, are able to generate clusters of proliferating chondrogenic cells that produce type II collagen and proteoglycans were obtained (Brun 1999).
- Bone Marrow Mesenchymal Stem are able to differentiate to several mesenchymal tissues including: bone, cartilage, connective tissue (Radice, 2000).

The in vitro reconstructed dermal substitutes obtained by culturing the patient own fibroblasts into a 3D Hyaluronan based scaffolds, are routinely used in the clinical practice in many hospital (Galassi et al 2000; Brun et al. 2000) for the treatment of skin lesion. Also autologous human chondrocytes cultured into the above described scaffolds are being used for the treatment of cartilage lesion.

Brun P, Dickinson SC, Zavan B, Cortivo R, Hollander AP, Abatangelo G.  
Characteristics of repair tissue in second-look and third-look biopsies from patients treated with engineered cartilage: relationship to symptomatology and time after implantation.  
*Arthritis Res Ther.* 2008;10(6):R132. Epub 2008 Nov 11.

Zavan B, Vindigni V, Vezzù K, Zorzato G, Luni C, Abatangelo G, Elvassore N, Cortivo R.  
Hyaluronan based porous nano-particles enriched with growth factors for the treatment of ulcers: a placebo-controlled study.  
*J Mater Sci Mater Med.* 2009 Jan;20(1):235-47. Epub 2008 Aug 30.

Zavan B, Abatangelo G, Mazzoleni F, Bassetto F, Cortivo R, Vindigni V.  
New 3D hyaluronan-based scaffold for in vitro reconstruction of the rat sciatic nerve.  
*Neurol Res.* 2008 Mar;30(2):190-6.

*Prof. Abatangelo*  
*Dept. Histology, Microbiology and Biomedical Technology*  
*University of Pavia, Italy*

Zavan B, Vindigni V, Lepidi S, Iacopetti I, Avruscio G, Abatangelo G, Cortivo R.

Neoarteries grown in vivo using a tissue-engineered hyaluronan-based scaffold. *FASEB J.* 2008 Aug;22(8):2853-61. Epub 2008 Apr 2.

Aguiari P, Leo S, Zavan B, Vindigni V, Rimessi A, Bianchi K, Franzin C, Cortivo R, Rossato M, Vettor R, Abatangelo G, Pozzan T, Pinton P, Rizzuto R. High glucose induces adipogenic differentiation of muscle-derived stem cells. *Proc Natl Acad Sci U S A.* 2008 Jan 29;105(4):1226-31. Epub 2008 Jan 22.

Hollander AP, Dickinson SC, Sims TJ, Brun P, Cortivo R, Kon E, Marcacci M, Zanasi S, Borriore A, De Luca C, Pavesio A, Soranzo C, Abatangelo G. Maturation of tissue engineered cartilage implanted in injured and osteoarthritic human knees. *Tissue Eng.* 2006 Jul;12(7):1787-98.

Lepidi S, Grego F, Vindigni V, Zavan B, Tonello C, Deriu GP, Abatangelo G, Cortivo R. Hyaluronan biodegradable scaffold for small-caliber artery grafting: preliminary results in an animal model. *Eur J Vasc Endovasc Surg.* 2006 Oct;32(4):411-7. Epub 2006 May 5.

Lepidi S, Abatangelo G, Vindigni V, Deriu GP, Zavan B, Tonello C, Cortivo R. In vivo regeneration of small-diameter (2 mm) arteries using a polymer scaffold. *FASEB J.* 2006 Jan;20(1):103-5. Epub 2005 Nov 15.

Ricciardolo FL, Caramori G, Ito K, Capelli A, Brun P, Abatangelo G, Papi A, Chung KF, Adcock I, Barnes PJ, Donner CF, Rossi A, Di Stefano A. Nitrosative stress in the bronchial mucosa of severe chronic obstructive pulmonary disease. *J Allergy Clin Immunol.* 2005 Nov;116(5):1028-35. Epub 2005 Sep 28.

Zavan B, Brun P, Vindigni V, Amadori A, Habeler W, Pontisso P, Montemurro D, Abatangelo G, Cortivo R. Extracellular matrix-enriched polymeric scaffolds as a substrate for hepatocyte cultures: in vitro and in vivo studies. *Biomaterials.* 2005 Dec;26(34):7038-45.

Tonello C, Vindigni V, Zavan B, Abatangelo S, Abatangelo G, Brun P, Cortivo R. In vitro reconstruction of an endothelialized skin substitute provided with a microcapillary network using biopolymer scaffolds. *FASEB J.* 2005 Sep;19(11):1546-8. Epub 2005 Jun 21.

Zavan B, Cortivo R, Tonello C, Abatangelo G. Gland cell cultures into 3D hyaluronan-based scaffolds. *J Mater Sci Mater Med.* 2003 Aug;14(8):727-9.

Brun P, Dickinson SC, Zavan B, Cortivo R, Hollander AP, Abatangelo G. Characteristics of repair tissue in second-look and third-look biopsies from patients treated with engineered cartilage: relationship to symptomatology and

*Prof. Abatangelo*  
*Dept. Histology, Microbiology and Biomedical Technology*  
*University of Pavia, Italy*

time after implantation.

Arthritis Res Ther. 2008;10(6):R132. Epub 2008 Nov 11.

Zavan B, Vindigni V, Vezzù K, Zorzato G, Luni C, Abatangelo G, Elvassore N, Cortivo R.

Hyaluronan based porous nano-particles enriched with growth factors for the treatment of ulcers: a placebo-controlled study.

J Mater Sci Mater Med. 2009 Jan;20(1):235-47. Epub 2008 Aug 30.

Zavan B, Abatangelo G, Mazzoleni F, Bassetto F, Cortivo R, Vindigni V.

New 3D hyaluronan-based scaffold for in vitro reconstruction of the rat sciatic nerve.

Neurol Res. 2008 Mar;30(2):190-6.

Zavan B, Vindigni V, Lepidi S, Iacopetti I, Avruscio G, Abatangelo G, Cortivo R.

Neoarteries grown in vivo using a tissue-engineered hyaluronan-based scaffold.

FASEB J. 2008 Aug;22(8):2853-61. Epub 2008 Apr 2.

Aguiari P, Leo S, Zavan B, Vindigni V, Rimessi A, Bianchi K, Franzin C, Cortivo R, Rossato M, Vettor R, Abatangelo G, Pozzan T, Pinton P, Rizzuto R.

High glucose induces adipogenic differentiation of muscle-derived stem cells.

Proc Natl Acad Sci U S A. 2008 Jan 29;105(4):1226-31. Epub 2008 Jan 22.

Hollander AP, Dickinson SC, Sims TJ, Brun P, Cortivo R, Kon E, Marcacci M, Zanasi S, Borriore A, De Luca C, Pavesio A, Soranzo C, Abatangelo G.

Maturation of tissue engineered cartilage implanted in injured and osteoarthritic human knees.

Tissue Eng. 2006 Jul;12(7):1787-98.

Lepidi S, Grego F, Vindigni V, Zavan B, Tonello C, Deriu GP, Abatangelo G, Cortivo R.

Hyaluronan biodegradable scaffold for small-caliber artery grafting: preliminary results in an animal model.

Eur J Vasc Endovasc Surg. 2006 Oct;32(4):411-7. Epub 2006 May 5.

Lepidi S, Abatangelo G, Vindigni V, Deriu GP, Zavan B, Tonello C, Cortivo R.

In vivo regeneration of small-diameter (2 mm) arteries using a polymer scaffold.

FASEB J. 2006 Jan;20(1):103-5. Epub 2005 Nov 15.

Ricciardolo FL, Caramori G, Ito K, Capelli A, Brun P, Abatangelo G, Papi A, Chung KF, Adcock I, Barnes PJ, Donner CF, Rossi A, Di Stefano A.

Nitrosative stress in the bronchial mucosa of severe chronic obstructive pulmonary disease.

J Allergy Clin Immunol. 2005 Nov;116(5):1028-35. Epub 2005 Sep 28.

Zavan B, Brun P, Vindigni V, Amadori A, Habeler W, Pontisso P, Montemurro D, Abatangelo G, Cortivo R.

Extracellular matrix-enriched polymeric scaffolds as a substrate for hepatocyte cultures: in vitro and in vivo studies.

Biomaterials. 2005 Dec;26(34):7038-45.



Tonello C, Vindigni V, Zavan B, Abatangelo S, Abatangelo G, Brun P, Cortivo R.

In vitro reconstruction of an endothelialized skin substitute provided with a microcapillary network using biopolymer scaffolds.

FASEB J. 2005 Sep;19(11):1546-8. Epub 2005 Jun 21.

Zavan B, Cortivo R, Tonello C, Abatangelo G.

Gland cell cultures into 3D hyaluronan-based scaffolds.

J Mater Sci Mater Med. 2003 Aug;14(8):727-9.

Brun P, Cortivo R, Zavan B, Vecchiato N, Abatangelo G.

In vitro reconstructed tissues on hyaluronan-based temporary scaffolding.

J Mater Sci Mater Med. 1999 Oct-Nov;10(10/11):683-8.

Giroto D, Urbani S, Brun P, Renier D, Barbucci R, Abatangelo G.

Tissue-specific gene expression in chondrocytes grown on three-dimensional hyaluronic acid scaffolds.

Biomaterials. 2003 Aug;24(19):3265-75.

Pavesio A, Abatangelo G, Borriero A, Brocchetta D, Hollander AP, Kon E, Torasso F, Zanasi S, Marcacci M.

Hyaluronan-based scaffolds (Hyalograft C) in the treatment of knee cartilage defects: preliminary clinical findings.

Novartis Found Symp. 2003;249:203-17; discussion 229-33, 234-8, 239-41. Review.

Brun P, Panfilò S, Daga Gordini D, Cortivo R, Abatangelo G.

The effect of hyaluronan on CD44-mediated survival of normal and hydroxyl radical-damaged chondrocytes.

Osteoarthritis Cartilage. 2003 Mar;11(3):208-16.

Tonello C, Zavan B, Cortivo R, Brun P, Panfilò S, Abatangelo G.

In vitro reconstruction of human dermal equivalent enriched with endothelial cells.

Biomaterials. 2003 Mar;24(7):1205-11.

Brun P, Grosso F, Galassi G, Radice M, Tonello C, Cortivo R, Abatangelo G.

Use of dermal-like tissue in the management of chronic and acute full-thickness cutaneous wounds.

Ostomy Wound Manage. 2000 Sep;46(9):44-8.

Brun P, Dickinson SC, Zavan B, Cortivo R, Hollander AP, Abatangelo G.

Characteristics of repair tissue in second-look and third-look biopsies from patients treated with engineered cartilage: relationship to symptomatology and time after implantation.

Arthritis Res Ther. 2008;10(6):R132. Epub 2008 Nov 11.

Zavan B, Vindigni V, Vezzù K, Zorzato G, Luni C, Abatangelo G, Elvassore N, Cortivo R.

Hyaluronan based porous nano-particles enriched with growth factors for the treatment of ulcers: a placebo-controlled study.

*Prof. Abatangelo*  
*Dept. Histology, Microbiology and Biomedical Technology*  
*University of Pavia, Italy*

J Mater Sci Mater Med. 2009 Jan;20(1):235-47. Epub 2008 Aug 30.

Zavan B, Abatangelo G, Mazzoleni F, Bassetto F, Cortivo R, Vindigni V.  
 New 3D hyaluronan-based scaffold for in vitro reconstruction of the rat sciatic nerve.

Neurol Res. 2008 Mar;30(2):190-6.

Zavan B, Vindigni V, Lepidi S, Iacopetti I, Avruscio G, Abatangelo G, Cortivo R.

Neoarteries grown in vivo using a tissue-engineered hyaluronan-based scaffold.  
 FASEB J. 2008 Aug;22(8):2853-61. Epub 2008 Apr 2.

Aguiari P, Leo S, Zavan B, Vindigni V, Rimessi A, Bianchi K, Franzin C,  
 Cortivo R, Rossato M, Vettor R, Abatangelo G, Pozzan T, Pinton P, Rizzuto R.  
 High glucose induces adipogenic differentiation of muscle-derived stem cells.  
 Proc Natl Acad Sci U S A. 2008 Jan 29;105(4):1226-31. Epub 2008 Jan 22.

Hollander AP, Dickinson SC, Sims TJ, Brun P, Cortivo R, Kon E, Marcacci M,  
 Zanasi S, Borriore A, De Luca C, Pavesio A, Soranzo C, Abatangelo G.  
 Maturation of tissue engineered cartilage implanted in injured and osteoarthritic human knees.

Tissue Eng. 2006 Jul;12(7):1787-98.

Lepidi S, Grego F, Vindigni V, Zavan B, Tonello C, Deriu GP, Abatangelo G, Cortivo R.

Hyaluronan biodegradable scaffold for small-caliber artery grafting: preliminary results in an animal model.

Eur J Vasc Endovasc Surg. 2006 Oct;32(4):411-7. Epub 2006 May 5.

Lepidi S, Abatangelo G, Vindigni V, Deriu GP, Zavan B, Tonello C, Cortivo R.  
 In vivo regeneration of small-diameter (2 mm) arteries using a polymer scaffold.  
 FASEB J. 2006 Jan;20(1):103-5. Epub 2005 Nov 15.

Ricciardolo FL, Caramori G, Ito K, Capelli A, Brun P, Abatangelo G, Papi A,  
 Chung KF, Adcock I, Barnes PJ, Donner CF, Rossi A, Di Stefano A.

Nitrosative stress in the bronchial mucosa of severe chronic obstructive pulmonary disease.

J Allergy Clin Immunol. 2005 Nov;116(5):1028-35. Epub 2005 Sep 28.

Zavan B, Brun P, Vindigni V, Amadori A, Habeler W, Pontisso P, Montemurro D, Abatangelo G, Cortivo R.

Extracellular matrix-enriched polymeric scaffolds as a substrate for hepatocyte cultures: in vitro and in vivo studies.

Biomaterials. 2005 Dec;26(34):7038-45.

Tonello C, Vindigni V, Zavan B, Abatangelo S, Abatangelo G, Brun P, Cortivo R.

In vitro reconstruction of an endothelialized skin substitute provided with a microcapillary network using biopolymer scaffolds.

FASEB J. 2005 Sep;19(11):1546-8. Epub 2005 Jun 21.

*Prof. Abatangelo*  
*Dept. Histology, Microbiology and Biomedical Technology*  
*University of Padoa, Italy*

Zavan B, Cortivo R, Tonello C, Abatangelo G.

Gland cell cultures into 3D hyaluronan-based scaffolds.

J Mater Sci Mater Med. 2003 Aug;14(8):727-9.

Brun P, Cortivo R, Zavan B, Vecchiato N, Abatangelo G.

In vitro reconstructed tissues on hyaluronan-based temporary scaffolding.

J Mater Sci Mater Med. 1999 Oct-Nov;10(10/11):683-8.

Giroto D, Urbani S, Brun P, Renier D, Barbucci R, Abatangelo G.

Tissue-specific gene expression in chondrocytes grown on three-dimensional hyaluronic acid scaffolds.

Biomaterials. 2003 Aug;24(19):3265-75.

Pavesio A, Abatangelo G, Borriore A, Brocchetta D, Hollander AP, Kon E, Torasso F, Zanasi S, Marcacci M.

Hyaluronan-based scaffolds (Hyalograft C) in the treatment of knee cartilage defects: preliminary clinical findings.

Novartis Found Symp. 2003;249:203-17; discussion 229-33, 234-8, 239-41. Review.

Brun P, Panfilo S, Daga Gordini D, Cortivo R, Abatangelo G.

The effect of hyaluronan on CD44-mediated survival of normal and hydroxyl radical-damaged chondrocytes.

Osteoarthritis Cartilage. 2003 Mar;11(3):208-16.

Tonello C, Zavan B, Cortivo R, Brun P, Panfilo S, Abatangelo G.

In vitro reconstruction of human dermal equivalent enriched with endothelial cells.

Biomaterials. 2003 Mar;24(7):1205-11.

Brun P, Grosso F, Galassi G, Radice M, Tonello C, Cortivo R, Abatangelo G.

Use of dermal-like tissue in the management of chronic and acute full-thickness cutaneous wounds.

Ostomy Wound Manage. 2000 Sep;46(9):44-8.

Galassi G, Brun P, Radice M, Cortivo R, Zanon GF, Genovese P, Abatangelo G.

In vitro reconstructed dermis implanted in human wounds: degradation studies of the HA-based supporting scaffold.

Biomaterials. 2000 Nov;21(21):2183-91.

Radice M, Brun P, Cortivo R, Scapinelli R, Battaliard C, Abatangelo G.

Hyaluronan-based biopolymers as delivery vehicles for bone-marrow-derived mesenchymal progenitors.

J Biomed Mater Res. 2000 May;50(2):101-9.

Benedetti L, Cortivo R, Berti T, Berti A, Pea F, Mazzo M, Moras M, Abatangelo G.

Biocompatibility and biodegradation of different hyaluronan derivatives (Hyaff) implanted in rats.

Biomaterials. 1993 Dec;14(15):1154-60.

Brun P, Abatangelo G, Radice M, Zacchi V, Guidolin D, Daga Gordini D, Cortivo

R.

Chondrocyte aggregation and reorganization into three-dimensional scaffolds.  
*J Biomed Mater Res.* 1999 Sep 5;46(3):337-46.

Chen WY, Abatangelo G.

Functions of hyaluronan in wound repair.  
*Wound Repair Regen.* 1999 Mar-Apr;7(2):79-89. Review.

Campoccia D, Doherty P, Radice M, Brun P, Abatangelo G, Williams DF.

Semisynthetic resorbable materials from hyaluronan esterification.  
*Biomaterials.* 1998 Dec;19(23):2101-27. Review.

Zacchi V, Soranzo C, Cortivo R, Radice M, Brun P, Abatangelo G.

In vitro engineering of human skin-like tissue.  
*J Biomed Mater Res.* 1998 May;40(2):187-94.

Abatangelo G, Barbucci R, Brun P, Lamponi S.

Biocompatibility and enzymatic degradation studies on sulphated hyaluronic acid derivatives.  
*Biomaterials.* 1997 Nov;18(21):1411-5.

Abatangelo G.

Hyaluronic acid delays or impedes reepithelialization?  
*J Burn Care Rehabil.* 1997 Nov-Dec;18(6):552. No abstract available.

Italiano G, Abatangelo G Jr, Calabrò A, Abatangelo G Sr, Zanoni R, O'Regan M, Passerini Glazel G.

Reconstructive surgery of the urethra: a pilot study in the rabbit on the use of hyaluronan benzyl ester (Hyaff-11) biodegradable grafts.  
*Urol Res.* 1997;25(2):137-42.

Cortivo R, Brun P, Cardarelli L, O'Regan M, Radice M, Abatangelo G.

Antioxidant effects of hyaluronan and its alpha-methyl-prednisolone derivative in chondrocyte and cartilage cultures.  
*Semin Arthritis Rheum.* 1996 Aug;26(1):492-501.

Cortivo R, Brun P, Rastrelli A, Abatangelo G.

In vitro studies on biocompatibility of hyaluronic acid esters.  
*Biomaterials.* 1991 Oct;12(8):727-30.

Brun P, De Galateo A, Camporese A, Cortivo R, Abatangelo G.

Analysis of hyaluronic acid in synovial fluid by reversed-phase liquid chromatography.  
*J Chromatogr.* 1990 Apr 6;526(2):530-4. No abstract available.

Cortivo R, De Galateo A, Castellani I, Brun P, Giro MG, Abatangelo G.

Hyaluronic acid promotes chick embryo fibroblast and chondroblast expression.  
*Cell Biol Int Rep.* 1990 Feb;14(2):111-22.

Abatangelo G, Botti P, Del Bue M, Gei G, Samson JC, Cortivo R, De Galateo A,

Martelli M.

Intraarticular sodium hyaluronate injections in the Pond-Nuki experimental model of osteoarthritis in dogs. I. Biochemical results.  
Clin Orthop Relat Res. 1989 Apr;(241):278-85.

Schiavinato A, Lini E, Guidolin D, Pezzoli G, Botti P, Martelli M, Cortivo R, De Galateo A, Abatangelo G.

Intraarticular sodium hyaluronate injections in the Pond-Nuki experimental model of osteoarthritis in dogs. II. Morphological findings.  
Clin Orthop Relat Res. 1989 Apr;(241):286-99.

Abatangelo G, Martelli M, Vecchia P.

Healing of hyaluronic acid-enriched wounds: histological observations.  
J Surg Res. 1983 Nov;35(5):410-6.

Abatangelo G, Cortivo R, Martelli M, Vecchia P.

Cell detachment mediated by hyaluronic acid.  
Exp Cell Res. 1982 Jan;137(1):73-8. No abstract available.